



Matthew H. Mead
Governor

State of Wyoming

Department of Workforce Services

DIVISION OF WORKERS' COMPENSATION

1510 East Pershing Boulevard, South Wing
Cheyenne, Wyoming 82002
<http://www.wyomingworkforce.org>



John Cox
Director
John Ysebaert
Deputy Director

PROVIDER BULLETIN

TOPIC: Facet Injections

Published by Wyoming Workers' Compensation Medical Case Management Unit and Effective on
June 28, 2017

September 14, 2017

The following is a clarification of diagnostic facet injections versus pain management injections:

- (1) Diagnostic facet injections must follow the guidelines as written on June 28, 2017.**
- (2) Once the pain generator(s) has been identified, multiple levels may be injected based on written pain treatment guidelines from the provider.**

Based on review of the current literature, the Medical Commission has determined the following guidelines will be implemented:

- Patient has a clinical picture of facet joint pain with cervical, thoracic or lumbar pain that can radiate to scapula, chest or buttock/ thigh, respectively. Tenderness to palpation over the suspected joint, pain increased by extension or extension and rotation of the spine. Neurologic examination is normal.
- Imaging studies prior to injection must demonstrate abnormality of the joint to be injected or demonstrate instability at the appropriate level on flexion/ extension films. The ideal imaging study, if available, would be CT-Nuclear SPECT, or if not available, Nuclear SPECT, injecting only levels that are "hot" on the imaging study.
- Facet injections will only be paid for by Workman's Compensation if appropriate follow-up is provided. This will be in the form of a written VAS kept by, and



signed by the patient. Local anesthetic and steroid of choice can be used. The VAS should include pre-injection pain level and post-injection pain level immediately after the injection and at each hour for 24 hours (while awake) as well as VAS at 48 and 72 hours. A diagram of pre-injection and post-injection pain location immediately after and one hour after the injection should be included.

- Only one level should be injected (uni or bi-lateral) at one session. Patient should be awake and no IV sedation or analgesics should be utilized. If multiple joints are pain generators, a series of blocks will be required, culminating in all symptomatic joints being injected simultaneously to completely relieve the patient's pain.
- Patients who have 80% relief of pain for the duration of the local anesthetic can be considered candidates for a Medial branch block and if that produces pain relief, then a Facet Rhizotomy can be performed.

